

Sixth degree Diophantine polynomial equation

Author: OLIVER COUTO

University of Waterloo

Email: matt345@celebrating-mathematics.com

Abstract

$$(abc)(ab+bc+ca)(a+b+c) = (pqr)(pq+pr+qr)(p+q+r) \text{ ----- (1)}$$

Historically, we note that finding a parametrization of degree six has not been easy. In the below paper the author has followed in the footsteps of below mentioned paper, ref. no. (1). In ref. no.(1), Ajai Choudhry on page 356 has parametrized equation: $m(abc)(ab+bc+ca)(a+b+c)=n(pqr)(pq+pr+qr)(p+q+r)$, where, $(m,n)=(3,32)$ using one parameter. In the below paper the author has parametrized the said equation for $(m,n)=(1,1)$, but with two parameters. The trick used is to split equation (1) into three parts, so as to balance the coefficients (m,n) & such that, $(m,n)=[(uvw),(xyz)]$. And $[(uvw),(xyz)]=(72,72)$, & hence they cancel each other out.

Consider the below equation:

$$(abc)(ab+bc+ca)(a+b+c)=(pqr)(pq+pr+qr)(p+q+r) \text{ ---(1)}$$

By the below method we arrive at a parametrization of equation (1) with two variables:

We split the above equation (1) as below:

$$u(a+b+c)=x(p+q+r) \text{ ----- (2)}$$

$$v(ab+bc+ca)=y(pq+qr+pr) \text{ ----- (3)}$$

$$w(abc)=z(pqr) \text{ ----- (4)}$$

We take: $(u,v,w)=(8,1,9)$ & $(x,y,z)=(9,4,2)$

$$8(a+b+c)=9(p+q+r) \text{ ----- (2)}$$

$$(ab+bc+ca)=4(pq+qr+pr) \text{ ----- (3)}$$

$$9abc=2pqr \text{ ----- (4)}$$

We note that: $(uvw)=(xyz)=72$

We take:

$$(a,b,c)=[(pu),(qv),(2r/9uv)] \text{ ----- (5)}$$

We notice that (4) is satisfied by (5) & Substituting (5) in (2) we get the value of variable 'r' as below:

$$r = \frac{9uv(p(8u - 9) + q(8v - 9))}{(81uv - 16)} \text{ ----- (6)}$$

We substitute value of 'r' from (6) in equation (3) & we simplify.

Hence we get a quadratic equation in (p,q) & after making the coefficient of (q^2) in it equal to zero at $(u = \frac{1}{18})$, we get the below values of variables (p, q).

$$p = (81v^2u^2 - 288vu^2 - 288uv^2 + 340uv - 18u - 18v + 64) \text{ ----- (7)}$$

$$q = 2u(144uv - 8u - 162v + 9) \text{ ----- (8)}$$

Substituting, $(u = \frac{1}{18})$, in (7) & (8) we get:

$$p = \frac{9(v^2 - 4)}{4} \text{ ----- (9)}$$

$$q = \frac{11(18v - 1)}{81} \text{ ----- (10)}$$

After substituting value of $(u = \frac{1}{18})$ & (p, q) from (9) & (10) in (6) we get the value of variable (r) as below:

$$r = \frac{11v(v-72)}{324} \text{ ----- (11)}$$

Since: $(a, b, c) = \left[(pu), (qv), \left(\frac{2r}{9uv} \right) \right]$, we get after substituting the values of $(u = \frac{1}{18})$ & (p, q, r) from (9), (10) & (11) we get:

$$a = \frac{(v^2 - 4)}{8}$$

$$b = \frac{11v(18v - 1)}{81}$$

$$c = \frac{11(v - 72)}{81}$$

After removing common factors we get, (a, b, c, p, q, r) , as below:

$$a = 81(v^2 - 4)$$

$$b = 88v(18v - 1)$$

$$c = 88(v - 72)$$

$$p = 1458(v^2 - 4)$$

$$q = 88(18v - 1)$$

$$r = 22v(v - 72)$$

We put, $(v = \frac{m}{n})$ & we get a parametrization in two variables & is shown below:

$$\mathbf{a = 81(m + 2n)(m - 2n)}$$

$$\mathbf{b = 88m(18m - n)}$$

$$\mathbf{c = 88n(m - 72n)}$$

$$\mathbf{p = 1458(m + 2n)(m - 2n)}$$

$$\mathbf{q = 88n(18m - n)}$$

$$\mathbf{r = 22m(m - 72n)}$$

For, $(m, n) = (3, 2)$, we get:

$$(a, b, c) = (567, -13728, 24816)$$

$$(p, q, r) = (10206, -9152, 9306)$$

I thank mathematician Dr. Ajai Choudhry, who readily sent (by email) the below parameterization (in three variables) of equation (1):

$$a = mu(m^2 + 1)(m^4 - m^2 + 1)(u + v)$$

$$b = mv(m^2 + 1)(m^4 - m^2 + 1)(u + v)$$

$$c = -m(m^6uv + u^2 + v^2 + uv)$$

$$p = u(m^2 + 1)(m^4 - m^2 + 1)(u + v)$$

$$q = v(m^2 + 1)(m^4 - m^2 + 1)(u + v)$$

$$r = -(m^6uv + m^6u^2 + m^6v^2 + uv)$$

Small numerical soln for above at, $(m, u, v) = (2, 2, 1)$:

$$(a, b, c, p, q, r) = (52, 26, -18, 26, 13, -30)$$

Acknowledgement:

The author is grateful to mathematician Seiji Tomita for his help & advice.

References

-
- 1) Ajai Choudhry, Journal of number theory, vol. 153, Aug 2015, pgs. 354-363
-

- 2) Dickson, L. E. History of the Theory of Numbers, Vol. 2: Diophantine Analysis, New York: Dover, 2005.
- 3) Oliver Couto, webpage, <http://www.celebrating-mathematics.com>
- 4) Seiji Tomita & Oliver Couto, Sums of three n th powers equal to a square. Bulletin of Mathematics & Statistics research, Vol.11, Issue #4, 2023 (Oct-Dec)
- 4) Construction of diagonal quartic & sextic surfaces with infinitely many rational points. Andrew Bremner, Ajai Choudhry, Maciej Ulas, Arxiv (2014), (#1402.4583)
- 5) Seiji Tomita, webpage <http://www.maroon.dti.ne.jp/fermat> Web site "Computation number theory"
- 6) Oliver Couto & Seiji Tomita, Generalized parametric solution to multiple sums of powers, Universal Journal of applied mathematics, Jan 2016, Volume 3(5),
Ramanujan lost notebook, Narosa publishing house, New Delhi, India <http://www.hrpub.org>
- 7) Oliver Couto & Seiji Tomita, Solution of equation of any degree ' n ' & $n = (2, 3, 4, 5 \text{ \& } 6)$, Journal "SOP transactions on applied mathematics" ISSN (Print): 2373-8472, July 2015.
- 8) Euler Leonhard, Opera Omnia books, 1984 publication.
- 9) Ajai Choudhry, Equal sums of three sixth powers, Rocky Mountain, Journal of mathematics, V30, 2000, Rocky mountain journal vol. 34, no.4.
- 10) Tito Piezas-Onlinebook on "Collection of algebraic identities", <http://sites.google.com/site/tpiezas>
- 11) Jaroslaw Wroblewski, Tables of Numerical solutions for degree 3,4,6,7. Website, www.math.uni.wroc.pl/~jwr/eslp
