

April 2020

Sums of Biquadratics with integer coefficients

Authors: *Oliver Couto*¹, *Seiji Tomita*²

1 University of Waterloo, Waterloo, Ontario, Canada
Email: samson@celebrating-mathematics.com

2 Computer Engineer, Tokyo software co. Inc., Japan.
Email: fermat@m15.alpha-net.ne.jp

Abstract

We consider two types of equations shown below:

$$ax^4 + by^4 = cz^4 + dw^4 \quad \text{----- (1)}$$

$$ax^4 + by^4 + cz^4 = dw^4 \quad \text{----- (2)}$$

Condition: product (abcd) not equal to zero

Existence of solution for Diophantine equation:

$ax^4 + by^4 = cz^4 + dw^4$ & $ax^4 + by^4 + cz^4 = dw^4$, are known if (abcd) is square number & product not equal to zero. So we are curious about whether above equation has a solution if (abcd) is not square number & product not equal to zero. In particular, when does this equation have infinitely many integer solutions? Bremner, A., & Choudhry, A., & Ulas [1] have showed the solution family of the similar equation ($ax^4 + by^4 + cz^4 + dw^4 = 0$) with infinitely many rational points using elliptic curve theory. We show other family of solution of this equation with infinitely many rational points. As a bonus we have considered an interesting case of equation ($ax^4 + by^4 = az^4 + bw^4$) in which the product of the co-efficients is a square number, but has been parameterized by using only algebraic methods without taking re-course to elliptic curve theory.

We have considered the below family of six equations:

$$(A) \quad p(X^4 + Y^4) = 2Z^4 + W^4$$

$$(B) \quad d^2(X^4) + (d - 2c)^2Y^4 = 2d(d - 2c)Z^4 + 4W^4$$

$$(C) \quad (d^2)(a^2)(X^4) + (-d + 2a)^2(a^2)(Y^4) + \\ 2d(-d + 2a)Z^4 = 4(a^2)W^4$$

$$(D) \quad pr^2(X^4) + 4p(p + r)^2Y^4 + r^3(Z^4) = r^2(p + r)W^4$$

$$(E) \quad a(p)^4 + b(q)^4 = c(r)^4 + b(d)^4$$

$$(F) \quad (-c^2 + 4ec - 2e^2)X^4 + (-c^2 + 4ec - 2e^2)Y^4 + \\ (2c - 2e)^2Z^4 = 2W^4$$

Case (A):

$$p(X^4 + Y^4) = 2Z^4 + W^4 \text{ --- (1)}$$

We use an identity,

$$p(t + 1)^4 + p(t)^4 = 2(t^2 + at + b)^2 + (ct^2 + dt + e)^2,$$

with,

$$p = 3 - 2d + \frac{(d)^2}{2},$$

$$e = 1, c = (-2 + d),$$

$$a = (-d + 3),$$

$$b = 1/2(2 - d)$$

So, we look for the integer solutions,

$$\left\{ Z^2 = t^2 + (-d + 3)t + 1 - \frac{d}{2}, \quad W^2 = (-2 + d)t^2 + dt + 1 \right\}$$

By parameterizing the second equation and substituting the result to first equation, then we obtain quartic equation below.

$$u^2 = (-2d + 4)k^4 + (-24 + 8d)k^3 + (32 - 4d)k^2 + (-8d^2 - 48 + 24d)k + (16 + 2d^3 - 4d^2)$$

This quartic equation has infinitely many rational solutions for

$$d = 6, 8, 18, 24, 26, 42, 48, 50 \text{ with } d \leq 50.$$

$$(d, p) = (6, 9), (8, 19), (18, 129), (24, 243), (26, 289), (42, 801), (48, 1059), (50, 1153).$$

Hence we can obtain infinitely many integer solutions for equation (1) where,

$$p = (9, 19, 129, 243, 289, 801) \text{ with } p < 1000$$

For $(d, p) = (6, 9)$ we get:

$$9x^4 + 9y^4 = 2z^4 + w^4$$

$$(x, y, z, w) = (41, 153, 223, 58)$$

Case(B):

$$d^2 a^2 X^4 + (-d + 2a)^2 a^2 Y^4 + 2d(-d + 2a)Z^4 = 4a^2 W^4$$

$$d^2 a^2 X^4 + (-d + 2a)^2 a^2 Y^4 + 2d(-d + 2a)Z^4 = 4a^2 W^4 \text{ --- (3)}$$

(a, d) are arbitrary.

We use an identity:

$$p(t + 1)^4 + q(t)^4 + r(at^2 + at)^2 = s(at^2 + dt + e)^2,$$

with $(p, q, r, s, e) =$

$$\left[\frac{1}{4}(sd^2), \frac{1}{4}(s(2a - d)^2), \frac{1}{2}(a^2)(sd(2a - d)), s, \frac{1}{2(d)} \right]$$

So, we look for the integer solutions:

$$\left\{ Z^2 = at^2 + at, \quad W^2 = at^2 + dt + \frac{1}{2d} \right\}$$

By parameterizing the second equation and substituting the result to first equation, then we obtain quartic equation below.

$$u^2 = (4a - 2d)k^4 + 2da^2$$

This quartic equation has infinitely many rational solutions for (a,d) with (a,d) ≤ 5 below.

$$(a, d) = (1,3), (1,4), (1,5), (3,2), (3,4), (4,1), (4,2), (4,3), (4,4), (4,5), (5,2).$$

Hence we can obtain infinitely many integer solutions for equation (3).

For (a,d)=(3,2) we have;

$$9x^4 + 36y^4 + 4z^4 = 9w^4$$

$$(x, y, z, w) = (1,12,6,17)$$

Case(C):

$$pr^2X^4 + 4p(p+r)^2Y^4 + r^3Z^4 = r^2(p+r)W^4$$

$$pr^2X^4 + 4p(p+r)^2Y^4 + r^3Z^4 = r^2(p+r)W^4 \text{---(4)}$$

$$pr^2X^4 + 4p(p+r)^2Y^4 + r^3Z^4 = r^2(p+r)W^4 \text{-----(4)}$$

p,r are arbitrary.

We use an identity $p^*(t+1)^4 + q^*(t)^4 + r^*(t^2 + at + 1)^2 = (p+r)^*(ct^2 + dt + 1)^2$,

with (a,c,d,q) = $(-2p/r, (2p+r)/r, 0, 4p(r^2 + p^2 + 2pr)/(r^2))$.

So, we look for the integer solutions $\{Z^2 = t^2 - \frac{2pt}{r} + 1, W^2 = (2p+r)t^2/r + 1\}$

By parameterizing the second equation and substituting the result to first equation, then we obtain quartic equation below.

$$u^2 = r^4k^4 + r(8pr^2 + 2r^3)k^2 + r(8pr^2 + 16p^2r)k + r(4p^2r + 8p^3 + r^3)$$

This quartic equation is birationally equivalent to an elliptic curve below.

$$\begin{aligned} Y^2 + (16r^5p + 32r^4p^2)Y \\ = X^3 + (8r^3p + 2r^4)X^2 + (-16r^6p^2 - 32r^5p^3 - 4r^8)X - 192r^9 * p^3 \\ - 32r^{10}p^2 - 256r^8p^4 - 32r^{11}p - 8r^{12} \end{aligned}$$

The corresponding point is:

$$P(X, Y) = (-8pr^3 - 2r^4, -16r^5p - 32r^4p^2).$$

Hence we get:

$$2P(X,Y) = (6r^4 + 4pr^3 + p^2r^2, 16r^5p - 10r^4p^2 + 16r^6 - p^3r^3).$$

This point P is of infinite order, and the multiples mP , $m = 2, 3, \dots$ give infinitely many points. This quartic equation (4) has infinitely many parametric solutions below.

$$m = 2:$$

$$X = -8r^2 - 8pr + p^2$$

$$Y = 4(2r + p)r$$

$$Z = 3p^2 - 4pr - 8r^2$$

$$W = p^2 + 12pr + 8r^2$$

$$m = 3:$$

$$X = (p^2 + 12pr + 8r^2)(-8r^2 - 8pr + p^2)(-64r^4 - 128r^3p - 112p^2r^2 - 48p^3r + p^4)$$

$$Y = 4(2r + p)(3p^2 - 4pr - 8r^2)r(p^2 + 12pr + 8r^2)^2$$

$$Z = (p^2 + 12pr + 8r^2)(5p^6 - 84p^5r - 728r^2p^4 - 832r^3p^3 + 576r^4p^2 + 1280r^5p + 512r^6)$$

$$W = (p^2 + 12pr + 8r^2)(p^6 + 100p^5r + 104r^2p^4 + 576r^3p^3 + 1856r^4p^2 + 1792r^5p + 512r^6)$$

For $(p,q)=(3,2)$ we have:

$$3x^4 + 75y^4 + 2z^4 = 5w^4$$

$$(x, y, z, w) = (71, 56, 26, 113)$$

Case(D):

$$(-c^2 + 4ec - 2e^2)X^4 + (-c^2 + 4ec - 2e^2)Y^4 + (2c - 2e)^2 * Z^4 = 2W^4.$$

$$(-c^2 + 4ec - 2e^2)X^4 + (-c^2 + 4ec - 2e^2)Y^4 + (2c - 2e)^2 * Z^4 = 2W^4..(7)$$

We use an identity:

$$p(t + 1)^4 + p(t)^4 + r(t^2 + at + b)^2 = 2(ct^2 + dt + e)^2,$$

with $\{a, b, d, p, r\} =$

$$\{1, 1/2(c - 2e)(c - e), c, -c^2 + 4ec - 2e^2, (2c - 2e)^2\}.$$

So, we look for the integer solutions:

$$\{Z^2 = t^2 + t + \frac{1}{2}(c - 2e)(c - e), W^2 = ct^2 + ct + e\}$$

By parameterizing the second equation and substituting the result to first equation, then we obtain quartic equation below.

$$u^2 = (-c^3 + 6c^2e - 8e^2c + 4e^3)k^4 + (8e^2c - 4c^2e)k^3 + (-28c^2e + 36e^2c + 6c^3 - 8e^3)k^2 + (8e^2c - 4c^2e)k - c^3 + 6c^2e - 8e^2c + 4e^3$$

This quartic equation has infinitely many rational solutions for $|(p,r)| < 100$ below.

$$(c, e) = (7,4), (9,6), (9,7), (9,10), (9,13)$$

Hence we can obtain infinitely many integer solutions for equation (7) where

$$(c, e) = (7,4), (9,6), (9,7), (9,10), (9,13)$$

For $(p,r)=(1,2)$ we get:

$$x^4 + 9y^4 + 2z^4 = 3w^4$$

$$(x, y, z, w) = (47,40,37,57)$$

Case (E):

$$a(p)^4 + b(q)^4 = c(r)^4 + d(s)^4$$

We take, $(c=a)$ & $(d=b)$ and we get:

$$a(p)^4 + b(q)^4 = a(r)^4 + b(s)^4 \text{ --- (1)}$$

Hence product of coefficients, $(abab) = (ab)^2$, equals a square

We multiply eqn. (1) by integer ten & we get

$$(10) * [a(p)^4 + b(q)^4] = (10) * [a(r)^4 + b(s)^4]$$

Hence we get:

$$(2)(5)(a)(p - r)(p + r)(p^2 + r^2) = (2)(5)(b)(s - q)(s + q)(s^2 + q^2)$$

Hence, we take, $(a, b) = ((s - q), (p - r))$

Let: $5(p + r) = 2(s + q)$

Applying the above conditions we get:

$$2(p^2 + r^2) = 5(s^2 + q^2) \text{ --- (2)}$$

Eqn.(1) has numerical solution at:

$$(p, q, r, s, a, b) = (11, 2, -7, 8, 1, 3) \text{ --- (3)}$$

We parameterize eqn. (2) at:

$$[(11 + 2t), (2 + 5kt), (-7 + 2kt), (8 + 5t)]$$

& we get after removing common factors:

$$p = 33k^2 - 8k + 17$$

$$q = 14k^2 + 40k - 6$$

$$r = 29k^2 + 16k + 21$$

$$s = 24k^2 - 20k - 16$$

And Since,

$$(a, b) = ((s - q), (p - r))$$

we have,

$$a = 19k^2 + 10k - 11$$

$$b = 31k^2 + 4k + 19$$

For $k=0$ we get:

$$(p, q, r, s) = (17, 2, 21, 16) \text{ \& } (a, b) = (-11, 19)$$

$$11(17)^4 + 19(16)^4 = 11(21)^4 + 19(6)^4$$

In the above parameterization we used above eqn. (3) numerical solution. Similarly there are other numerical solutions listed below that can be used to arrive at more parameteric forms for eqn. (1). This can be done by finding suitable relationship for the eqn.

$$m(p + r) = n(s + q)$$

Where, (m, n) are integers.

Numerical examples for equation (1):

$$(a, b, p, q, r, s) = (3, 5, 6, 1, 4, 5)$$

$$(a, b, p, q, r, s) = (1, 33, 24, 13, 9, 14)$$

$$(a, b, p, q, r, s) = (33, 59, 96, 17, 22, 83)$$

$$(a, b, p, q, r, s) = (1, 11, 21, 12, 1, 14)$$

$$(a, b, p, q, r, s) = (15, 31, 7, 20, 24, 5)$$

$$(a, b, p, q, r, s) = (1, 3, 4, 1, 2, 3)$$

$$(a, b, p, q, r, s) = (11, 27, 11, 12, 16, 1)$$

$$(a, b, p, q, r, s) = (4, 9, 1, 14, 17, 6)$$

Case(F):

$$d^2X^4 + (d - 2c)^2Y^4 = 2d(d - 2c)Z^4 + 4W^4.$$

$$d^2X^4 + (d - 2c)^2Y^4 = 2d(d - 2c)Z^4 + 4W^4..(8)$$

We use an identity:

$$p(t + 1)^4 + q(t)^4 = r(t^2 + at + b)^2 + s(ct^2 + dt + e)^2,$$

$$p = d^2, q = (d - 2c)^2, r = 2d(d - 2c), s = (2)^2$$

with $\{a, b, e, p, q, r\} =$

$$\{1, 0, 1/2(d), 1/4(sd^2), 1/4(sd^2) - scd + sc^2,$$

$$1/2(sd^2) - scd\}.$$

So, we look for the integer solutions:

$$\left\{ Z^2 = t^2 + t, \quad W^2 = ct^2 + dt + \frac{1}{2}(d) \right\}$$

By parameterizing the first equation and substituting the result to second equation, then we obtain quartic equation below.

$$u^2 = (4c - 2d)k^4 + 2d$$

Quartic curve is transformed to the elliptic curve E.

$$x = (4c - 2d)k^2, y = (4c - 2d)uk.$$

$$E: y^2 = x^3 + 2(4c - 2d)dx$$

If rank of 'E' is greater than zero, then 'E' has infinitely many rational solutions.

Hence we can obtain infinitely many integer solutions for equation (8) if rank of E is greater than zero.

$$|c| < 5, d < 10$$

$$[c, d]rank [p, q, r, s] [x, y, z, w]$$

We give below numerical solutions to the equation:

$$p(x)^4 + q(y)^4 = r(z)^4 + s(w)^4$$

$$[c, d]rank [p, q, r, s] [x, y, z, w]$$

For (c,d)=(-4,1) & (p,q,r,s)=(1,81,18,4) we get:

$$x^4 + 81y^4 = 18z^4 + 4w^4$$

$$(x, y, z, w) = (9, 1, 3, 6)$$

See Table below:

Table (1)

[-4, 1] [1]	[1, 81, 18, 4]	[9, 1, 3, 6]
[-4, 1] [1]	[1, 81, 18, 4]	[9801, 2209, 4653, 5106]
[-4, 5] [2]	[25, 169, 130, 4]	[9, 1, 3, 14]
[-4, 5] [2]	[25, 169, 130, 4]	[49, 9, 21, 74]
[-4, 5] [2]	[25, 169, 130, 4]	[14161, 8649, 11067, 3886]
[-4, 8] [1]	[16, 64, 64, 1]	[12769, 7056, 9492, 15934]

[-1, 2] [1]	[1, 4, 4, 1]	[9, 4, 6, 7]
[-1, 2] [1]	[1, 4, 4, 1]	[12769, 7056, 9492, 7967]
[-1, 8] [1]	[16, 25, 40, 1]	[81, 16, 36, 158]
[0, 3] [1]	[9, 9, 18, 4]	[49, 1, 7, 60]
[0, 7] [1]	[49, 49, 98, 4]	[6241, 2209, 3713, 10920]
[1, 3] [1]	[9, 1, 6, 4]	[9, 1, 3, 11]
[1, 3] [1]	[9, 1, 6, 4]	[361, 625, 475, 13]
[1, 3] [1]	[9, 1, 6, 4]	[201601, 27889, 74983, 246121]
[1, 4] [1]	[4, 1, 4, 1]	[169, 1, 13, 239]

Table (Contd.):

$$p(x)^4 + q(y)^4 = r(z)^4 + s(w)^4$$

[c, d] rank [p, q, r, s] [x,y,z,w]

Table (2):

[1, 8] [1]	[16, 9, 24, 1]	[49, 16, 28, 94]
[1, 8] [1]	[16, 9, 24, 1]	[3721, 1089, 2013, 7199]
[1, 9] [2]	[81, 49, 126, 4]	[9, 1, 3, 19]
[1, 9] [2]	[81, 49, 126, 4]	[25, 1, 5, 53]
[1, 9] [2]	[81, 49, 126, 4]	[1521, 529, 897, 3071]
[1, 9] [2]	[81, 49, 126, 4]	[3481, 2401, 2891, 5861]
[1, 9] [2]	[81, 49, 126, 4]	[6241, 5329, 5767, 8711]

[1, 9] [2]	[81, 49, 126, 4]	[42849, 47089, 44919, 22391]
[1, 9] [2]	[81, 49, 126, 4]	[72361, 37249, 51917, 136771]
[4, 1] [1]	[1, 49, -14, 4]	[2209, 1, 47, 1562]
[4, 2] [1]	[1, 9, -6, 1]	[1, 4, 2, 7]
[4, 2] [1]	[1, 9, -6, 1]	[121, 9, 33, 122]
[4, 2] [1]	[1, 9, -6, 1]	[169, 225625, 6175, 390794]
[4, 2] [1]	[1, 9, -6, 1]	[2209, 784, 1316, 2593]
[4, 3] [1]	[9, 25, -30, 4]	[1521, 529, 897, 2042]
[4, 5] [1]	[25, 9, -30, 4]	[529, 1521, 897, 2042]
[4, 6] [1]	[9, 1, -6, 1]	[4, 1, 2, 7]
[4, 6] [1]	[9, 1, -6, 1]	[9, 121, 33, 122]
[4, 6] [1]	[9, 1, -6, 1]	[225625, 169, 6175, 390794]

Note: Above Tables may facilitate further research in equation (1).

References:

[1], Bremner Andrew , Ajai Choudhry, & Ulas M, (2014), Constructions of diagonal quartic and sextic surfaces with infinitely many rational points, International Journal of Number Theory, 10(7), 1675-1698.

[2] A.S.Elsehanes & J. Jalind, Diophantine equation $(x^4+2y^3=z^4+4w^4)$, math computations ,75,2005,pages 935-940.

[3] Andrew Bremner & M. Ulas, International journal of number theory, vol. 7, number. 8 (2011), pages.2081-2090

[4] Ajai Choudhry, Diophantine equations, Journal of mathematics, Rocky mountain journal vol. 34, no.4, winter (2004), page1261-1298.

[5] Oliver Couto, personal webpage, <http://www.celebrating-mathematics.com> Web site on “Number theory Math”

[6] J.H. Silverman: a) Rational points on elliptic curves, b) Arithmetic of elliptic Curves, Springer 1992

- [7] Crenome's & Steinwatkins tables for elliptic curves
- [8] Elliptic curves tables: [http://www. Imfdb.org](http://www.Imfdb.org)
- [9] Published math paper, Oliver Couto & Seiji Tomita,), Sum of three biquadratics, Universal Journal of applied mathematics , Vol. (4)(1), pages 22-31,2016,
- [10] Ramanujan lost notebook, Narosa publishing house, 1990, New Delhi, India
- [11] Seiji Tomita, personal webpage<http://www.maroon.dti.ne.jp/fermat> Web site on "Computation number theory"
- [12] Equal Sums of Quartics (In Context with the Richmond Equation) ($ax^2+by^4+cz^4+dw^4=0$). Universal Journal of Applied Mathematics Vol. 8(2): 13 - 29, 2020, DOI:10.13189/ ujam.2020.08020 <http://www.hrpub.org>
- [13] Published math paper, Oliver Couto & Seiji Tomita, Generalized parametric solution to multiple sums of powers, Universal Journal of applied mathematics, In press, Jan 2016, Volume 3(5), <http://www.hrpub.org>
- [14] Euler Leonhard, Opera Omnia books, 1984 publication.
- [15] Published math paper, Oliver Couto & Seiji Tomita, Solution of Polynomial Equation of any degree 'n' with Special Emphasis For n= (2, 3, 4, 5 & 6), Journal- -"SOP transactions on applied-mathematics"ISSN(Print):2373-8472, July,2015. <http://www.scipublish.com/journals/am>
- [16] Tito Piezas-Online collection algebraic identities <http://sites.google.com/site/tpiezas>
- [17] Published math paper, Oliver Couto, Taxicab Equations for power (2, 3, 4 & 5), Journal, International Math Forum, Hikari ltd., Vol.9,Jan,2014.n0.12, pages 561-577. <http://www.m-hikari.com/imf.html>
- [18] Jaroslaw Wroblewski , Tables of Numerical solutions for degree three, four, six, Seven & nine, website www.math.uni.wroc.pl/~jwr/eslp
- [19] N.D. Elkies on sum of three quartics equal to a quartic, Math computation 51, (184), 1988, pages 825-835
- [20] L. E. Dickson, history of the theory of numbers, Vol. II, (Diophantine analysis), Chelsea publication, year 2000
-